



**NATIONAL HEMOPHILIA FOUNDATION**  
for all bleeding and clotting disorders

## Physical Therapy Practice Guidelines for Persons with Bleeding Disorders: Joint Bleeds

The following practice guidelines were developed through the consensus of the therapists that work with patients with bleeding disorders and edited by the National Hemophilia Foundation's Physical Therapy Working Group. The information contained in the practice guidelines is not intended in any way to be used as primary medical advice or to replace medical advice. They are intended to guide the physical therapist caring for individuals with bleeding disorders in the important factors and elements of quality care.

### Signs and Symptoms

- |   |   |
|---|---|
| <ul style="list-style-type: none"> <li>• Decreased ROM</li> <li>• Pain</li> <li>• Swelling/edema</li> <li>• Heat/erythema</li> <li>• Ecchymosis</li> <li>• Muscle guarding of affected joint</li> <li>• Muscle spasm</li> </ul> | <ul style="list-style-type: none"> <li>• Disuse of affected joint</li> <li>• Gait deviations: decreased weight bearing, antalgic</li> <li>• Decreased functional mobility/ activities of daily living</li> <li>• Pediatrics: decline in developmental milestones/motor pattern (i.e. resort to crawling)</li> </ul> |
|---|---|

### Differential Diagnosis

- |   |   |
|---|---|
| <ul style="list-style-type: none"> <li>• Trauma: Fracture, Avulsion, Dislocation, Ligament Injury, Cartilage injury</li> <li>• Synovitis</li> <li>• Pseudotumor</li> <li>• Bony or soft tissue tumor</li> <li>• Cyst</li> <li>• Sprain/strain</li> <li>• Neuroma</li> </ul> | <ul style="list-style-type: none"> <li>• Internal derangement</li> <li>• Infection ( intra or extra-articular)Inflammatory arthritis</li> <li>• Muscle bleeds</li> <li>• Complex Regional Pain Syndrome</li> <li>• Musculoskeletal ultrasound (if available) can be used to determine if a bleed is present in the joint space. Musculoskeletal ultrasound (if available) can be used to determine if excess fluid &amp;/or soft tissue proliferation is present in the joint space.</li> </ul> |
|---|---|

### PT Intervention

**Acute Stage:** pain present at rest and with all activity

- Factor replacement per medical team orders (consider prophylaxis prior to PT)
- RICE (ice for 10-20 minutes every 1-2 hours: see [Cryotherapy Guidelines](#))
- Immobilize / Splint
- NWB with appropriate assistive device

**Subacute Stage:** ADLs do not increase pain from baseline (will have some baseline pain)

- Factor replacement per medical team orders (consider prophylaxis prior to PT)
- Continue RICE prn for pain and after exertion • Splinting, resting splint for protection at night, weaning splint as able with pain as guide
- Wean assistive device and begin progressive weight bearing
- Activity modification to avoid pain
- Pain-free gentle AROM of affected joint
- Pain-free progressive strengthening of affected and unaffected joint

This material is provided for your general information only. NHF does not give medical advice or engage in the practice of medicine. NHF under no circumstances recommends particular treatment for specific individuals and in all cases recommends that you consult your physician or local treatment center before pursuing any course of treatment.

## Chronic Stage: pain-free with ADLs

- Factor replacement per medical team orders (consider prophylaxis prior to PT)
- Dynamic splinting to increase ROM prn
- Progressive active/passive ROM to affected joint
- Progressive Strengthening: Open chain, closed chain, resistive band and proprioceptive training to affected joint
- Gentle joint mobilizations to increase ROM
- Modified functional activities and sports to avoid aggravation of symptoms
- Orthotics or possible casting for chronic issues
- Conditioning to address other areas of impairment
- Education regarding precautions to prevent re-injury/re-bleeds or injury to other joints

## Other Treatment Considerations

- TENS
- Electrical Stimulation
- Relaxation Techniques for muscle guarding
- Kinesiology tape
- Cryotherapy (see [Cryotherapy Guidelines](#))
- Elastic stockinette
- Joint support for return to sports
- Appropriate treatment if associated sprain/injury
- Myofascial release
- Further consultation with medical team
- Work with hematologist for adequate factor coverage
- Further imaging (plain radiographs to determine joint status, MRI to determine the presence of chronic synovitis)
- Treatment duration will vary based on individual needs and may require a longer duration than those identified for individuals without bleeding disorders.

## Precautions/Contraindications

Inhibitor patients: Please contact the physical therapist at your regional Hemophilia Treatment Center (HTC) for more information on these complex patients

- Avoid aggressive exercise and activity
- Increasing activity too early
- Monitor for nerve compression
- Use of heat modalities including ultrasound: (refer to MASAC Guideline #130 at [www.hemophilia.org](http://www.hemophilia.org))
- No return to sport/activity until pain-free full ROM and strength

This material is provided for your general information only. NHF does not give medical advice or engage in the practice of medicine. NHF under no circumstances recommends particular treatment for specific individuals and in all cases recommends that you consult your physician or local treatment center before pursuing any course of treatment.

## References

- Auerswald G, Dolan G, Duffy A, Hermans C, Jiménez-Yuste V, Ljung R, ... Šalek S Z. Pain and Pain Management in Haemophilia. *Blood Coagulation & Fibrinolysis*, 2016, 27(8), 845–854. doi.org/10.1097/MBC.0000000000000571
- Carcao M, Hilliard P, Escobar M A, Solimeno L, Mahlangu J and Santagostino E. Optimising Musculoskeletal Care for Patients with Haemophilia. *Eur J Haematol*, 2015, 95: 11–21. doi:10.1111/ejh.12581
- Di Minno M, Pasta G, Airaldi S, Zaottini F, Storino A, Cimino E, and Martinoli C. Ultrasound for Early Detection of Joint Disease in Patients with Hemophilic Arthropathy. *Journal of Clinical Medicine*, 2017, 6(8), 77. doi:10.3390/jcm6080077
- Hermans C, De Moerloose P, Fischer K, Holstein K, Klamroth R, Lambert T, Lavigne-Lissalde G, Perez R, Richards M, Dolan G and on the behalf of the European Haemophilia Therapy Standardization Board (2011), Management of acute haemarthrosis in haemophilia A without inhibitors: literature review, European survey and recommendations. *Haemophilia*, 2017: 383–392. doi:10.1111/j.1365-2516.2010.02449.x
- Kuijlaars IAR, Timmer MA, de Kleijn P, Pisters MF, Fischer K. Monitoring Joint Health in Haemophilia: Factors Associated with Deterioration. *Haemophilia*. 2017; 00:1-7. doi.org/10.1111/hae.13327
- Lobet S, Hermans C, and Lambert C. Optimal Management of Hemophilic Arthropathy and Hematomas. *Journal of Blood Medicine*, 2014, 5, 207–218. doi.org/10.2147/JBM.S50644
- Rodriguez-Merchan EC. Articular Bleeding in Hemophilia. *Cardiovascular & Hematological Disorders-Drug Targets*, volume 16, issue 1, pages 21-24, yr2016, issn 1871-529X/2212-4063, doi 10.2174/1871529X16666160613114506
- Strike K, Mulder K, Michael R. Exercise for Haemophilia. *Cochrane Database of Systematic Reviews* 2016, Issue 12. Art. No.: CD011180. DOI: 10.1002/14651858.CD011180.pub2
- Thornburg CD, Duncan NA. Treatment Adherence in Hemophilia. *Patient Preference and Adherence* 2017: 11 1677–1686. doi:10.2147/PPA.S139851
- Wyseure T, Mosnier LO, von Drygalski A. Advances and Challenges in Hemophilic Arthropathy, In *Seminars in Hematology*, Volume 53, Issue 1, 2016, Pages 10-19, ISSN 0037-1963, doi.10.1053/j.seminhematol.2015.10.005

Please visit NHF's website @ [www.hemophilia.org](http://www.hemophilia.org) for HTC contact information and PT references. Additional information is also available through NHF @ 1 800 42-HANDI.

This material is provided for your general information only. NHF does not give medical advice or engage in the practice of medicine. NHF under no circumstances recommends particular treatment for specific individuals and in all cases recommends that you consult your physician or local treatment center before pursuing any course of treatment.